

- How the bladder works
- What goes wrong?
- What investigations need to be done?
- Who can help?
- Advice
- Medication
- Equipment
- Other help
- Useful contacts

Continence in MSA

Bladder problems are very common in people with MSA and are often amongst the first symptoms of the illness.

Even if bladder problems cannot be cured, they can be managed and this factsheet explains how.

How the bladder works

Urine is produced in the kidneys and passes down tubes called ureters into the bladder.

The bladder, in which urine is stored, is balloon shaped. It is situated low in the abdomen behind the pubic bone. From the bladder the urine leaves the body through another tube, the urethra. The valves or sphincters at the base of the bladder makes it watertight and stop urine leakage.

The bladder can hold about 1 pint (500ml) of urine but a reduction in its capacity is a common feature of many bladder disorders. The first sensation of needing to go to the toilet happens when the bladder is about two thirds full.

It used to be thought that the bladder was merely a waterproof bag but in the last 15 years it has been shown that the walls of the bladder are made up of several layers, some of which have the sole function of generating nerve signals to convey information about the bladder's contents to the spinal cord and the brain. If it is socially appropriate to go to the toilet, the brain sends messages down the spinal cord to relax the sphincter (the valve which opens the bladder) and to contract the bladder wall muscle (the detrusor) so that urine is passed from the urethra.

It is normal to pass urine between four to seven times during the day and once at night. This may sound straightforward, but it is a very complex process and needs the bladder, the brain



and all the nerves between them to be working in co-ordination. Being able to control this process is what is required for continence.

What goes wrong?

Bladder problems may occur for reasons such as prostatic outflow obstruction in men and stress incontinence in women, so it is important to consider other possible causes before putting them all down to MSA. This information however, concentrates on the common problems arising in MSA.

MSA seems to attack the neurological controlling centres of the bladder in quite a selective way. This is why bladder symptoms occur as an early feature of the disease, and why they may become particularly troublesome over ensuing years. People with MSA get a combination of an overactive bladder, incomplete emptying and sphincter weakness. Some of the potential problems are:

- **Urgency** - needing to go to the toilet suddenly and quickly. This is due to the bladder contracting of its own accord - ~~bladder overactivity~~ bladder overactivity. If you cannot reach the toilet in time you may suffer urge incontinence. However, urge incontinence is a common symptom in the general population, not just in people with MSA.
- **Frequency** - going to the toilet a lot (more than eight times in 24 hours) but only passing small amounts of urine. This can either be due to the bladder's overactivity or because you are not emptying it completely.
- **Double voiding** - needing to pass urine again very soon after just doing so indicating your bladder was not completely empty the first time, sometimes accompanied by a feeling of incomplete emptying.
- **Urine infections** - especially if you have incomplete bladder emptying. Urine infections often make people feel very unwell and can make other symptoms, such as postural hypotension (a sudden fall in blood pressure), worse. The early use of antibiotics to treat urine infections is very important for people with MSA.
- **Constipation** - adding to problems of poor bladder function.

Problems with going to the toilet can also be affected by other MSA symptoms:

- Passing lots of urine during the night; this usually happens to people who also have postural hypotension as a symptom.
- Accidental leakage, because of difficulty or slowness getting to the toilet or adjusting clothing in time.

What investigations might be done?

Trying to find out exactly what the problem may be is an important starting point for getting appropriate treatment. When you see a nurse or a doctor they will ask you questions about your general health, how much you drink, the colour and smell of your urine and your bladder problems, including whether you have had any accidental leakage.

You may be asked to produce a sample of urine to test for a range of different things, including signs of infection. The sample can be sent to a laboratory for detailed testing or tested immediately with special urine testing sticks. The sticks give quite a good indication as to whether or not the urine is infected, but a more reliable result comes from the laboratory. If it is felt necessary to grow the germs in urine on a special plate to test for antibiotic sensitivity, it may take several days to get a result.

Measuring the volume and frequency of passing urine can help determine what your exact problem is. You may be asked to keep a diary for a few days, recording everything you drink and how often you pass urine and even measuring the volumes.

The other investigation that is very useful is checking how much urine is left in your bladder after you have passed urine, using either a small ultrasound machine or a catheter. Any urine left is called the post-micturition residual urine volume.

All these tests can be done at your home or at your GP's surgery. If more detailed testing is needed, you may be referred for urodynamics at the hospital. Urodynamics are a range of procedures that test how well your bladder fills and empties.

Urological treatments for several common bladder problems may involve surgery, but operations in people with MSA are rarely the solution. It is not uncommon for people with MSA to have had some urological surgery, which may not have helped very much, before the neurological diagnosis is recognised. Further surgery in this case is probably inadvisable.

Who can help?

District nurses and Community Matrons can give advice on continence issues and will refer on to Continence advisors as appropriate

Continence advisors are healthcare professionals who have specialist training and experience in managing bladder and bowel problems. Continence advisors work in hospitals, health centres and in the community. Many accept self-referrals over the telephone, although some may ask that your GP write to them. The Bladder and Bowel Community helpline will have details of your nearest advisor (see [Helpful Contacts](#) at the end of this factsheet).

Your GP or specialist will also be able to help.

The Trust's MSA Nurse Specialists are also available to discuss bladder problems and treatment (see [Helpful Contacts](#)).

Advice

Sensible advice about fluid, toilet habits or diet is often very useful in preventing problems and managing symptoms.

Maintaining a healthy bladder means drinking plenty of fluid (about eight large cups or glasses each day). Drinking too little fluid can irritate your bladder and cause problems. There is some evidence that drinking cranberry juice may reduce urinary tract infections slightly, and continence advisors recommend reducing your intake of caffeine and fizzy drinks.

Be comfortable on the toilet, especially in public lavatories. Ladies should sit rather than hover (carry some wipes with you to do this) and men should use cubicles which provide privacy.

Some people find bending forward and gently pressing or slow firm tapping over the bladder at the end of the flow helps to squeeze out any urine left in the bladder.

Choosing clothes that are easy to undo or take off, for example Velcro fastenings rather than zips, can give you valuable time which may prevent accidental leakage.

Medication

Medication, called anti-muscarinic drugs (also known as anti-cholinergics), allows the bladder to relax and fill closer to capacity before needing to empty. These can reduce the symptoms of urgency and frequency.

DDAVP (desmopressin) is a hormone that prevents urine production for several hours after it has been taken. It can be useful if taken at night as it can stop the need to get up to go to the toilet, which improves sleep.

It comes in a tablet and nasal spray but it must only be used once a day and is not recommended for those over 65 years as it can cause water intoxication. It may be useful for people who have postural hypotension as a symptom.

Research suggests that botox injections into the bladder can be helpful for people with bladder problems caused by neurological disease. Speak to your MSA Nurse Specialist or Consultant about further information on this.

Regular or daily laxatives may be required if constipation is contributing to bladder problems.

Equipment

An occupational therapist can help to make it easier to use the toilet at home. Adjusting the height of the toilet, adding grab rails or even creating a downstairs toilet are all worth thinking about.

Community nurses can help provide urinals or commodes to make toileting easier.

There is a wide range of continence pads now available with varying absorbency suitable for day or night use by men and women and which can be either disposable or washable. Some are available on prescription and the continence nurse or community nurse can arrange to supply them to you.

Some men find urinary sheaths useful. They fit over the penis like a condom and attach to a urine collection bag via a tube. These are not usually used for the whole day, as they can cause skin irritation, but they can be useful when out and about or overnight.

Catheters are thin plastic tubes that can be passed into the bladder to drain it and can either be ~~intermittent~~ or ~~indwelling~~ (left in for some days or weeks).

Intermittent catheters are inserted into the urethra to reach the bladder, the urine is drained (which only takes a few minutes) and then the catheter is removed. A continence advisor will teach you how to do this for yourself or possibly show your carer how to do it for you. Intermittent catheterisation is a very useful means of improving bladder control if not emptying is a significant problem.

Indwelling catheters are also inserted via the urethra or through the abdomen (a supra pubic catheter). The catheter drains urine either into a drainage bag or has an attached valve that allows the bladder to be drained at regular intervals.

Although no one likes the idea of using a catheter many people are surprised at how easy they are to use and how they can relieve bladder problems.

Both types of catheter can be discreet and can give people more freedom from needing to use the toilet. However, for many people the decision to use any kind of catheter needs to be given plenty of thought and discussion.

Other help

'Can't wait' card: This credit card sized card can be shown discreetly to gain easy access to toilets when you are away from home. It is free and copies are available from the MSA Trust office or from the Bladder and Bowel Community.

The National Key Scheme: Initiated by RADAR, this offers independent access for disabled people into over 4,000 locked public toilets around Britain. Ideally all accessible toilets should be kept unlocked, but the scheme is used where it is necessary to lock the toilets to maintain

their cleanliness and to protect them from vandalism and misuse. Keys can be purchased from Disability Rights UK - see [Helpful Contacts](#)

Social Services: In some cases financial assistance towards laundry equipment, e.g. a washing machine, or to have clothes and bedding laundered, may be available. A social worker or benefits advisor can provide more details.

Sexual activity and continence: Bladder problems don't mean the end of intimate or sexual relationships. Some continence advisors are experienced in offering advice or treatment to enable sexual activity to be maintained. They recognise that for some people this is a very important part of life so will not be surprised or embarrassed if you want to talk about this with them. During a visit, they may even ask you directly if you are experiencing any problems with sexual function so you may want to prepare your reply. The MSA Trust also has a factsheet about sex and relationships which is available by contacting the office.

Useful contacts:

The Bladder and Bowel Community

The Foundation provides information, advice and expertise to anyone with bladder and bowel problems.

Telephone: 0800 031 5412 (Medical Helpline)

By post: Bladder & Bowel Community, 7 The Court, Holywell Business Park, Northfield Road, Southam CV47 0FS

Website: www.bladderandbowel.org

Disability Rights UK - National key scheme for locked toilets.

Telephone: 020 7250 8181

By post: Disability Rights UK, Ground Floor, CAN Mezzanine, 49-51 East Road, London N1 6AH

Website: www.disabilityrightsuk.org

Disabled Living Foundation - This is a national charity providing equipment advice and information for disabled people.

Telephone: 0300 999 0004 (Helpline)

By post: Disabled Living Foundation, Unit 1, 34 Chatfield Road, Wandsworth, London SW11 3SE

Website: www.dlf.org.uk

The Trust's contact details:

MSA Trust, 51 St Olav's Court, City Business Centre, Lower Road, London SE16 2XB

T: 0333 323 4591 | E: support@msatrust.org.uk | W: www.msatrust.org.uk

The MSA Trust Nurse Specialists:

Samantha Pavey (South East & East England): T: 0203 371 0003 | E: samantha.pavey@msatrust.org.uk

Katie Rigg (Scotland, Ireland and North England): T: 01434 381 932 | E: katie.rigg@msatrust.org.uk

Jill Lyons (Wales & South West England): T: 01934 316 119 | E: jill.lyons@msatrust.org.uk

Bowel Management - The factsheet titled **Bowel Management and MSAqis** available from the Trust office.

REVISION DATE: 09/17 | REVIEW DATE: 09/19 | VERSION: 1.3



Disclaimer

We have taken every care to ensure the accuracy of the information contained in this publication. It is produced independently, is not influenced by sponsors and is free from endorsement. The information should not be used as a substitute for the advice of appropriately qualified professionals, if in any doubt please seek advice from your doctor or legal professional.

References for this information sheet are available by contacting support@msatrust.org.uk.

Feedback

Your feedback helps us ensure we are delivering information to the highest standard. If you have any comments or suggestions please complete a short survey by following the links from our website: www.msatrust.org.uk or by contacting us at support@msatrust.org.uk.