

- What is a Gastrostomy feeding tube? Page 1
- When to start thinking about Gastrostomy feeding tube Page 1
- Can anyone have a Gastrostomy feeding tube? Page 2
- The benefits of Gastrostomy tube feeding Page 2
- What a Gastrostomy feeding tube does not do Page 3
- How is the Gastrostomy feeding tube fitted? Page 3
- What are the risks? Page 3
- What happens next? Page 4
- What can I put down the tube? Page 5
- Recuperation Page 5
- What else do I need to know? Page 5
- What happens if I decide not to have a feeding tube? Page 5

Gastrostomy Tube Feeding

This factsheet has been developed to help people with MSA understand why tube feeding may be something to consider as a supplement or alternative to taking food by mouth. It describes the process of having a gastrostomy tube fitted and explains the risks. A PEG tube is the most common way of inserting a gastrostomy feeding tube.

What is a Gastrostomy feeding tube?

A gastrostomy tube is a thin tube that comes into the stomach through a small hole made in the abdomen. This allows liquid feeds, fluids and medications to go directly into your stomach without the need for you to swallow it.

When to start thinking about a Gastrostomy feeding tube

Your Doctor, Speech and Language Therapist or Nurse Specialist (this could be an MSA, Parkinson's or Neurology Nurse) may discuss with you the possibility of having a tube for feeding if they recognise that you are showing some early signs of changes with your swallow.

Tube Feeding

It is good to start considering this early so you have time to think about what is involved in having a –gastrostomy tube inserted, whether this is something you would want and time to ask any questions you may have about it.

It is best to think about a gastrostomy tube before considerable weight loss has occurred and before you are at risk of chest infection due to aspiration (inhalation) of food into the airway. Before making a decision, you may want to discuss it with your family and other professionals such as, your Speech and Language Therapist and Doctor.

Can anyone have a Gastrostomy feeding tube?

There may be some people with MSA for whom siting a gastrostomy feeding tube is not an option due to other health considerations or because they are too frail and unwell generally for the procedure to be done safely. The medical team assessing you for a gastrostomy feeding tube will only undertake the procedure if they feel it will be of benefit to you and not cause greater harm.

Some people may not be able to have a feeding tube fitted because their stomach sits too high under the ribcage or the intestines over-ride the stomach getting between the stomach and abdominal wall. This prevents the feeding tube being passed through the stomach and abdominal wall as it would damage the intestines.

For most people with MSA having a gastrostomy tube fitted is something that can be planned for and you can choose whether you think this is a good option for you.

Sometimes a gastrostomy tube is fitted when someone is already in hospital having been particularly unwell and so is done as a more urgent procedure to assist their recovery. Wherever possible it is best to have a chance to think about whether you would want a gastrostomy feeding tube. There is no right or wrong answer. What is important is getting a chance to ask any questions that you may have and the information you need to make the decision that is right for you.

Continuing to eat and drink

Many people with MSA continue to take some foods and/or drinks by mouth after a gastrostomy tube has been fitted. You may even enjoy them more because you can eat just small amounts and only the foods you like, without having to worry about getting enough to eat. Your Speech and Language Therapist will advise you about what is safe for you and what types of food and drink are suitable.

The benefits of gastrostomy tube feeding

Difficulty in swallowing can become distressing for all concerned. Gastrostomy tube feeding can remove this anxiety, which often accompanies meals and allows you to get all the nutrition you need from a specially prepared liquid feed.

Sometimes the effort and the time it takes to eat a full meal reduces the pleasure we normally get from eating and drinking. A gastrostomy tube can be used to ‘top-up’ your nutrition if after 20 minutes or so you have had enough of eating. It is also a good way to increase your fluid intake, especially if you need to take extra fluid to maintain a good blood pressure, or maintain good urinary flow through a urinary catheter. Being able to increase your fluid intake generally will also help reduce fatigue, reduce constipation, keep your airway secretions loose and reduce urine infections.

Gastrostomy tube feeding may reduce the risk of chest infections, which can result from small amounts of food and/or fluid entering the airway during swallowing. However, the

Tube Feeding

muscle at the top of the stomach may be less efficient in people with MSA so food can back up into the food pipe and still go down the airway. To reduce this risk it is suggested you sit as upright as possible whilst eating and during a gastrostomy tube feed. Remain sitting up for 1 to 2 hours if possible after a feed. Also, chest infections can occur from saliva going down the airway, these will not be prevented by having a gastrostomy tube.

For most people with MSA having gastrostomy tube feeding provides an improved sense of wellbeing as your energy levels increase. If you have lost a lot of weight your dietitian may prescribe a high calorie feed to help you regain some of it. After this, the type and amount of feed will be adjusted to help keep your weight at the right level for you.

Many people with MSA have trouble with their bowels. The reasons for this include weakened pelvic floor muscles and poor fibre and/or fluid intake associated with swallowing difficulties. It can take a while for the bowels to get back to a normal pattern and your Dietitian may advise the use of a special feed with extra fibre if constipation is a problem.

A gastrostomy tube can be used to have your medications too. Not all medications can be put down a gastrostomy tube but most can and the pharmacist can advise you on the appropriate preparations of the medications you take. This can be particularly helpful when acutely unwell as at these times swallowing can be more difficult than usual and you may have extra medication to take.

What gastrostomy tube feeding does not do

Being fed by gastrostomy tube does not stop the progression of MSA or stop all risks of aspiration pneumonia. As already mentioned, saliva and reflux of stomach contents can still be aspirated into the lungs causing pneumonia.

How is the gastrostomy tube fitted?

The 2 most usual ways for a gastrostomy tube to be fitted are via a PEG or a RIG

PEG stands for: **P**ercutaneous - through the skin; **E**ndoscopic - using an endoscope (a thin flexible tube with a light on the end) **G**astrostomy - opening into the stomach

RIG stands for **R**adiologically (under Xray) **I**nserted **G**astrostomy – opening in the stomach

A short stay in hospital of 2-5 days is usually required to site a gastrostomy, ensure the site is clean, teach you how to manage it and establish what feed / fluids you will have through the tube and when.

Difference between **PEG** and **RIG**

A **PEG** tube is usually fitted under sedation. You will not be unconscious, but you will have an injection that makes you relax and provides pain relief to minimise discomfort. For a PEG to be fitted you will need to be able to lie fairly flat.

The endoscope is passed down the throat and is swallowed into the stomach. This does not impair your breathing and although it may be uncomfortable it is not painful.

The light of the endoscope highlights the inside of the stomach wall to identify the best place to pass the gastrostomy tube through. The gastrostomy tube is then passed down and comes through the stomach wall and out onto the surface skin of the abdomen through a small incision.

For some people lying down or having sedation may not be a good option for them, then a **RIG** is carried out. In this case you can stay sitting up and do not need sedation. You will



Multiple System Atrophy Trust | 128B Business Design Centre, 52 Upper Street, London, N1 0QH

Telephone 0333 323 4591 | Website www.msatrust.org.uk

Registered Charity Number 1137652 (England & Wales) and SC044535 (Scotland)

Tube Feeding

have been given barium a day or so before to show up the stomach on Xray. A small incision is made in the skin of the abdomen through to the stomach and the gastrostomy tube is inserted through this.

Whether the tube is fitted by PEG or RIG procedure, you will need a short stay in hospital, at least overnight, sometimes for a few days.

- A qualified Nurse will remain with you throughout the procedure.
- In both procedures the gastrostomy tube is held on the inside of the stomach by a small disc or water filled balloon on the end of the tube to prevent it from falling out. A small flange on the tube on the outside of the abdomen slides up the tube to hold it securely against the skin surface of the abdomen.
- If the gastrostomy tube is held in place by a disc inside the stomach this tube will not need to be routinely replaced. If the tube becomes damaged or needs replacing this will be done in hospital by the team who insert the tubes.
- If the tube is held inside the stomach by a balloon filled with water (the water is changed weekly through a valve on the side of the tube), this tube is usually changed every 12 weeks by a Nurse who specialises in tube care. They will visit you at home to do this.

What are the risks?

- With the PEG procedure a sore throat after the endoscope part of the procedure is common but should resolve after a few hours
- There is a small risk of damage to crowned teeth or dental bridgework when passing the endoscope in the PEG procedure.
- There is a small risk of an infection occurring during the procedure, most commonly when the tube comes out onto the abdominal skin surface. An antibiotic will be given to reduce the risk and treat any infection
- There is a small risk of bleeding where the tube passes through into the stomach
- There is a very small risk of significant problems with gastrostomy tube insertion, the team doing the procedure will discuss this with you.

What happens next?

Whilst in hospital you will be shown how to look after the tube, how to clean the area around the tube and how to use the tube for food, fluids and taking medication.

What can I put down the tube?

Through the tube you can take:

- Your food - There are many types of specially formulated liquid feeds that can go down the gastrostomy tube and your Dietician will determine which is the best for your needs. The Dietician will arrange for the feeds and supplies to be prescribed by your GP and delivered directly to you
- Medication in liquid form
- Water
- It may be possible for other fluids to be taken but this must be discussed with your Dietitian first as some liquids can corrode the tube.

You can choose to use the gastrostomy tube as much as you wish, and you can still eat and drink what you can safely manage and enjoy. We strongly recommend that you follow the advice of your Speech and Language Therapist regarding this.



Multiple System Atrophy Trust | 128B Business Design Centre, 52 Upper Street, London, N1 0QH

Telephone 0333 323 4591 | Website www.msatrust.org.uk

Registered Charity Number 1137652 (England & Wales) and SC044535 (Scotland)

Tube Feeding

People often feel much better after the gastrostomy tube is in place. Mealtimes can be less of a struggle and it can assist effective nourishment and hydration. Weight loss may become less of an issue. You should feel less tired, hungry and thirsty than before the gastrostomy tube was inserted.

Initially, you will be given water through the gastrostomy tube and the dietitian will work out an individual regime for you to ensure you have the correct calorific intake and nutritional balance.

Recuperation

The stomach and abdomen will heal in five to seven days. Any discomfort can be treated with medication.

You, your carer or family will be taught:

- How to care for the skin around the tube
- Signs and symptoms of infection
- What to do if the tube is pulled out
- Signs and symptoms of tube blockage
- What feeds to use and how often
- How to hide the tube under clothing
- When normal activities can be resumed.

What else do I need to know?

You will be able to take a shower soon after the gastrostomy tube is inserted and after two weeks you can take a bath and go swimming

You can go out and will find the gastrostomy tube can be easily tucked away under clothing

It is wise to ensure the tube is curled up and securely fastened to your abdomen to reduce the risk of it being caught and pulled when getting dressed or moving about.

What happens if I decide not to have a feeding tube?

Overtime with MSA eating and drinking becomes more difficult for most people, so without a tube you will lose weight and be less well hydrated. As this happens, gradually your body adjusts, and you become less thirsty and hungry.

The Speech and Language Therapist and Dietician will help you and advise about modifying food and fluid intake at this time.

The Trust's contact details

We have MSA Health Care Specialists that support people affected by MSA in the UK and Ireland. If you would like to find the MSA Health Care Specialist for your area, contact us on the details below or use the interactive map here – <https://www.msatrust.org.uk/support-for-you/hcps/>.

T: 0333 323 4591

E: support@msatrust.org.uk

W: www.msatrust.org.uk



Multiple System Atrophy Trust | 128B Business Design Centre, 52 Upper Street, London, N1 0QH

Telephone 0333 323 4591 | Website www.msatrust.org.uk

Registered Charity Number 1137652 (England & Wales) and SC044535 (Scotland)

Revision date: 11/25 | Review date: 11/28 | Version: 1.6

Disclaimer:

We have taken every care to ensure the accuracy of the information contained in this publication. However, the information should not be used as a substitute for the advice from appropriately qualified professionals. Speak to your doctor, qualified health care professional or legal professional before taking any action. Please note that personal views and opinions expressed are not necessarily endorsed by the Trust. The resource is produced independently, is not influenced by sponsors and is free from endorsement.

References for this information sheet are available by contacting support@msatrust.org.uk.

Your feedback helps us ensure we are delivering information to the highest standard. If you have any comments or suggestions, please contact us at support@msatrust.org.uk.



Multiple System Atrophy Trust | 128B Business Design Centre, 52 Upper Street, London, N1 0QH
Telephone 0333 323 4591 | Website www.msatrust.org.uk
Registered Charity Number 1137652 (England & Wales) and SC044535 (Scotland)